



WELCOME TO YOUR JOURNEY WITH NPLATE[®]



Inside you'll find helpful information on what you can expect from your Nplate[®] treatment as you are getting started.

OVER 15 YEARS of experience with **>500,000 PATIENTS** served globally

Indication:

Nplate[®] is a prescription medicine used to treat low blood platelet counts (thrombocytopenia) in adults with immune thrombocytopenia (ITP) when certain medicines or surgery to remove your spleen have not worked well enough.

Nplate[®] is not for use in people with a precancerous condition called myelodysplastic syndrome (MDS), or low platelet count caused by any condition other than immune thrombocytopenia (ITP). Nplate[®] is only used if your low platelet count and medical condition increase your risk of bleeding. Nplate[®] is used to try to keep your platelet count about 50,000 per microliter in order to lower the risk for bleeding. Nplate[®] is not used to make your platelet count normal. It is not known if Nplate[®] works or if it is safe in people under the age of 1.

Please see Important Safety Information throughout and on page 7.

WELCOME TO NPLATE®



WHAT IS NPLATE®?

Nplate®, a once-weekly in office injection, is a platelet booster that works with your body to create more platelets. Nplate® is used after steroids and does not work by suppressing the immune system. Instead, it increases the activities of the cells that naturally produce platelets.

A normal platelet count ranges from 150,000-450,000 per microliter of blood. However, **your healthcare team does not look for your treatment to get you back to this range.** Instead, **they will look to get your platelets between 50,000-200,000 to reduce the risk of bleeding.** Make sure you have a conversation with your healthcare team on platelet count goals.

If you'd like to learn more about ITP, visit [Nplate.com/ITP](https://www.nplate.com/ITP)

Important Safety Information

What is the most important information I should know about Nplate®?

Nplate® can cause serious side effects, including:

- **Worsening of a precancerous blood condition to a blood cancer (leukemia):** Nplate® is not for use in people with a precancerous condition called myelodysplastic syndromes (MDS), or for any condition other than immune thrombocytopenia (ITP). If you have MDS and receive Nplate®, your MDS condition may worsen and become an acute leukemia. If MDS worsens to become acute leukemia you may die sooner from the acute leukemia.

Please see additional Important Safety Information on page 7.

It's important to take an active role in your treatment journey.

The best way to do this is by asking questions and talking with your healthcare team. Throughout this brochure, we included key questions to help you start the conversation. See questions below.



QUESTIONS FOR YOUR HEALTHCARE TEAM:

- Ask how Nplate® works and how it is different than other treatments
- Discuss platelet count goals and understand what may be realistic for you




HOW FAST COULD NPLATE® WORK FOR PATIENTS?


During a clinical study, 93% of patients achieved a platelet response with Nplate®. Of those patients, some responded as early as 1 week, but it could take several weeks or more to find the right dose to achieve platelet stability. **It's important to give Nplate® time to work, but discuss what is best for your treatment journey with your healthcare team.**

As you can see from the image below, for some people it could take longer than 1-2 weeks. Your healthcare team can adjust your dosage as needed to find the dose to stabilize your platelets.

Of the Nplate® patients with a platelet response during clinical trials, **how soon did they see results?**



Responded in weeks 1-2



Responded in weeks 3-6



Responded in weeks 7-10



It's important to know that even if your platelet counts are considered stable (at or above 50,000), it's normal if they fluctuate weekly. Your healthcare team will determine if your Nplate® dosage needs to be adjusted.

In a clinical study of 75 patients, 6 out of 10 people kept their platelet counts at or above 50,000 per microliter for 11 months or more. This was the primary objective of the study and the majority of people reached a platelet count of at least 50,000 after 1 to 3 weekly doses.

Important Safety Information

• Higher risk for blood clots:

- You may have a higher risk of getting a blood clot if your platelet count becomes high during treatment with Nplate®. You may have severe complications or die from some forms of blood clots, such as clots that spread to the lungs or that cause heart attacks or strokes.
- If you have a chronic liver disease, you may get blood clots in the veins of your liver. This may affect your liver function.

Please see additional Important Safety Information on page 7.

TREATMENT-FREE REMISSION MAY BE POSSIBLE WITH NPLATE®

What does this mean? Some patients on Nplate® could stop treatment and still maintain at a stable platelet count.



About 1 in 3 adults went into treatment-free remission with Nplate®

This means that during the clinical study, their platelet counts **reached and stayed at or above 50,000** so that they were able to **stop any ITP treatment for at least 6 months**.



About half of those adults achieved treatment-free remission by about 6 months with Nplate®

However, the time it took to reach treatment-free remission varied from less than 2 months to more than 13 months.



QUESTIONS FOR YOUR HEALTHCARE TEAM:

- Discuss timing expectations around when you may see your platelets stabilize
- If you aren't seeing your platelets stabilize, ask if a dose adjustment might be right for you

 **Nplate**[®]
romiplostim injection

WHAT YOU MAY EXPECT AT YOUR WEEKLY APPOINTMENT

1

Check your current platelet count

After you check in for your appointment, you will have your blood drawn to get your CBC. This will help determine what dose of Nplate[®] is needed.

Note: Your healthcare team may also ask you to get your CBC checked a day or so before your appointment vs the day of. This may take place at another facility (e.g., a Labcorp or Quest). Either way is okay.

2

Your healthcare team will review your CBC results and adjust your Nplate[®] dosage based on your platelet count that week

3

The pharmacist will prepare your Nplate[®] dose

This may be done by the Pharmacist, Nurse or another Healthcare Provider depending upon your office.

4

Your Nplate[®] dose is administered

Important Safety Information

• What are the possible side effects of Nplate[®]?

- Nplate[®] may cause serious side effects. See **“What is the most important information I should know about Nplate[®]?”**
- The most common side effects of Nplate[®] in adults include headache, joint pain, dizziness, trouble sleeping, muscle tenderness or weakness, pain in arms and legs, stomach (abdomen) pain, shoulder pain, indigestion, tingling or numbness in hands and feet, bronchitis, inflammation of the sinuses (sinusitis), vomiting, diarrhea, upper respiratory tract infection, cough, nausea, and pain in mouth and throat (oropharyngeal pain).

Please see additional Important Safety Information on page 7.

WITH NPLATE[®]:



Continue to eat your favorite foods on your schedule

Nplate[®] can be taken with or without food and there are no dietary restrictions



No need to remember to take a daily pill

Nplate[®] is a once-weekly injection



Continue to take your other medications

Nplate[®] has no known drug interactions (e.g., statins, multivitamins)



No liver monitoring requirements with Nplate[®]



QUESTIONS FOR YOUR HEALTHCARE TEAM:

- If your appointment location is too far, discuss if there is an option to receive your Nplate[®] injection closer to home
- If your schedule interferes with your current appointment time, ask about changing it to another day that works for you



WELCOME

TREATMENT
EXPECTATIONS

YOUR
APPOINTMENT

PERSONALIZED
DOSING

SUPPORT

SAFETY

UNDERSTANDING PERSONALIZED DOSING WITH NPLATE®

You are unique, and how you respond to Nplate® may differ from someone else on treatment. **With Nplate®, there may be an opportunity to find the dose that works for you to stabilize your platelets!**

Your healthcare team will monitor your platelet count and adjust your Nplate® dosage accordingly. Each week, your Nplate® dosage may be adjusted if needed, starting at 1mcg/kg up to 10mcg/kg to try to find the right dose that keeps your platelet levels stable.

A stable dose is the dose at which platelet counts are at or above 50,000 per microliter for 4 or more weeks without dose adjustments.

Note: When starting on any ITP therapy, you should expect to get weekly CBCs to monitor your platelets. **If stability is achieved, platelet count monitoring can be reduced from weekly to monthly.**

Important Safety Information

- People who take Nplate® may have an increased risk of developing new or worsening changes in the bone marrow called "increased reticulin." These changes may improve if you stop taking Nplate®. Your healthcare provider may need to check your bone marrow for this problem during treatment with Nplate®.
- These are not all the possible side effects of Nplate®. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. For more information, ask your healthcare provider or pharmacist.
- If you have any questions about this information, be sure to discuss with your doctor. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information on page 7.

Take control of your treatment journey and keep track of your platelet count each week by using our **PLATELET TRACKER!**



Scan the QR code
to download!




QUESTIONS FOR YOUR HEALTHCARE TEAM:

- It's normal for your platelet count to fluctuate each week. Discuss what platelet count range is considered acceptable
- Discuss your platelet count goals and how your dosage may be adjusted accordingly
- If your platelet count is stable, ask about reducing the frequency of how often your platelet counts are checked



THE MAJORITY OF PATIENTS PAY \$0 OUT-OF-POCKET FOR NPLATE®*



*Based on an analysis from January 2022 to December 2023 for the Commercial and Medicare patients claims combined. Data based on IQVIA LAAD claims data. Oral OOP percent is a weighted average of claims from eltrombopag, avatrombopag, and fostamatinib disodium hexahydrate.

Important Safety Information

How will I receive Nplate®?

Nplate® is given as an injection under the skin (subcutaneous) one time each week by your healthcare provider. Your healthcare provider will check your platelet count every week and change your dose of Nplate® as needed. This will continue until your healthcare provider decides that your dose of Nplate® can stay the same. After that, you will need to get blood tests every month. When you stop receiving Nplate®, you will need blood tests for at least 2 weeks to check if your platelet count drops too low.

Please see additional Important Safety Information on page 7.

AMGEN® Support⁺

We're right here, right when you need us



The Amgen SupportPlus Co-Pay Program

If you have private or commercial insurance that you get from your employer or buy directly from a health insurance company, you may be eligible to **pay as little as \$0*** for your prescription.

Check your eligibility and sign up today at www.AmgenSupportPlus.com/copay

*Eligibility criteria and program maximums apply. See AmgenSupportPlus.com/copay for full Terms and Conditions.



Financial Support

We know every patient has unique needs. And we're here to provide financial support information and resources, regardless of your current financial situation or type of insurance you have.

Visit AmgenSupportPlus.com to learn more.



Amgen Nurse Partners[‡] can offer support to help you on your treatment journey.

Get connected with an Amgen Nurse Partner by enrolling in Amgen SupportPlus:

Download and complete the [Amgen SupportPlus Enrollment Form](#) found on AmgenSupportPlus.com and fax to **866-218-4777**

Call **866-264-2778** to enroll by phone

[‡]Amgen Nurse Partners are only available to patients that are prescribed certain Amgen products. They are not part of your treatment team and do not provide medical advice, nursing, or case management services. Amgen Nurse Partners will not inject patients with Amgen medications. Patients should always consult their healthcare provider regarding medical decisions or treatment concerns.



Scan the QR code for more information and helpful resources or **visit nplate.com/support**.



IMPORTANT SAFETY INFORMATION

What is the most important information I should know about Nplate®?

Nplate® can cause serious side effects, including:

- **Worsening of a precancerous blood condition to a blood cancer (leukemia):** Nplate® is not for use in people with a precancerous condition called myelodysplastic syndromes (MDS), or for any condition other than immune thrombocytopenia (ITP). If you have MDS and receive Nplate®, your MDS condition may worsen and become an acute leukemia. If MDS worsens to become acute leukemia you may die sooner from the acute leukemia.
- **Higher risk for blood clots:**
 - You may have a higher risk of getting a blood clot if your platelet count becomes high during treatment with Nplate®. You may have severe complications or die from some forms of blood clots, such as clots that spread to the lungs or that cause heart attacks or strokes.
 - If you have a chronic liver disease, you may get blood clots in the veins of your liver. This may affect your liver function.
- **What are the possible side effects of Nplate®?**
 - Nplate® may cause serious side effects. See **“What is the most important information I should know about Nplate®?”**
 - The most common side effects of Nplate® in adults include headache, joint pain, dizziness, trouble sleeping, muscle tenderness or weakness, pain in arms and legs, stomach (abdomen) pain, shoulder pain, indigestion, tingling or numbness in hands and feet, bronchitis, inflammation of the sinuses (sinusitis), vomiting, diarrhea, upper respiratory tract infection, cough, nausea, and pain in mouth and throat (oropharyngeal pain).
- People who take Nplate® may have an increased risk of developing new or worsening changes in the bone marrow called “increased reticulin.” These changes may improve if you stop taking Nplate®. Your healthcare provider may need to check your bone marrow for this problem during treatment with Nplate®.
- These are not all the possible side effects of Nplate®. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. For more information, ask your healthcare provider or pharmacist.

- If you have any questions about this information, be sure to discuss with your doctor. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

How will I receive Nplate®?

Nplate® is given as an injection under the skin (subcutaneous) one time each week by your healthcare provider. Your healthcare provider will check your platelet count every week and change your dose of Nplate® as needed. This will continue until your healthcare provider decides that your dose of Nplate® can stay the same. After that, you will need to get blood tests every month. When you stop receiving Nplate®, you will need blood tests for at least 2 weeks to check if your platelet count drops too low.

Indication:

Nplate® is a prescription medicine used to treat low blood platelet counts (thrombocytopenia) in adults with immune thrombocytopenia (ITP) when certain medicines or surgery to remove your spleen have not worked well enough.

Nplate® is not for use in people with a precancerous condition called myelodysplastic syndrome (MDS), or low platelet count caused by any condition other than immune thrombocytopenia (ITP). Nplate® is only used if your low platelet count and medical condition increase your risk of bleeding. Nplate® is used to try to keep your platelet count about 50,000 per microliter in order to lower the risk for bleeding. Nplate® is not used to make your platelet count normal. It is not known if Nplate® works or if it is safe in people under the age of 1.

Please see [Prescribing Information and Medication Guide](#) for more information about Nplate® or visit Nplate.com.

CBC, complete blood count; ITP, immune thrombocytopenia; OOP, out-of-pocket.